

Tel: 705-721-3344 Fax: 705-721-5522

BOWEL PREPARATION FOR COLONOSCOPY

Bi-Peglyte PREP

PLEASE ENSURE YOU HAVE READ THROUGH THE ENTIRE DOCUMENT REGARDING COLONOSCOPY (SEPARATE DOCUMENT) THAT PERTAINS TO APPOINTMENT TIME, PROCEDURE DATE, LOCATION, INSTRUCTIONS FOR MEDICATIONS, ETC.

STEP 1:

PURCHASE AT ANY PHARMACY:

- Bi-Peglyte BOWEL PREP
- Package include 2 Sachets and a blister pack with 3 Bisacodyl tabs



STEP 2:

- Day Before Procedure:
 - You must have only clear fluids for the entire day and evening.
 - NO DAIRY, NO SOLID FOOD, NO ALCOHOL.
 - You may have clear broth, apple juices, carbonated clear beverages, popsicles, Jell-O, coffee or tea, sports drinks (Powerade, Gatorade, etc.)
 - YOU MUST AVOID ANY RED, PURPLE, OR ORANGE DRINKS AS THESE COLOURS CAN INTEREFERE WITH THE PROCEDURE.

STEP 3:

- Day Before Procedure:
 - Prepare the Peg/Electrolyte solution by dissolving the entire contents of one sachet in 1L (32 ounces) of water. Mix rapidly for 2-3 minutes until a clear solution forms.
 - o Repeat steps above to make another 1L solution using second sachet.
 - o Refrigerate the 2L to improve taste. DO NOT DRINK SOLUTION YET.

STEP 4:

- Day Before Procedure:
 - At 3PM take 3 tablets of Bisacodyl tablets (total of 15 mg) with water.
 This medication does not prevent you from going to work.
 - At 7PM Drink 1L of Peg/Electrolyte solution within 1 hour. Continue to drink additional clear fluids as it is important to keep hydrated.



15 Gallie Court, Suite 306, Barrie, ON

Tel: 705-721-3344 Fax: 705-721-5522

STEP 5:

- Morning of Procedure:
 - 5 hours before scheduled arrival time Drink another 1L of Peg/Electrolyte solution within 1 hour.
 - **O DO NOT EAT OR DRINK ANYTHING ELSE ASIDES FROM WATER.**
 - You should take your morning medications with sips of water. See other document for instructions on which medications to take.

DO NOT DRINK ANYTHING AT ALL FOR 2 HOURS PRIOR TO YOUR ARRIVAL.

PLEASE DO YOUR BEST TO ARRIVE AT YOUR APPOINTMENT ON TIME.